

## Comment

### HIV: consensus indicators are needed for concurrency

Concurrent sexual partnerships are hypothesised to be the distinguishing factor behind the severe HIV epidemics in sub-Saharan Africa,<sup>1,2</sup> where 71% of the new HIV infections globally in 2008 occurred.<sup>3</sup> Although concurrency is informally recognised as overlapping sexual partnerships,<sup>4</sup> a precise definition and agreed indicator have not emerged. Several studies of sexual behaviour suggest that concurrent partnerships are more common and accepted in sub-Saharan Africa than elsewhere,<sup>5,6</sup> and mathematical models show that concurrent partnerships could increase the size and growth rate of HIV epidemics.<sup>7</sup> Unfortunately, the empirical evidence is weakened by the proliferation of measures for concurrency used in different studies,<sup>8</sup> some of which do not even capture whether or not partnerships overlap.<sup>9</sup> In part, the various definitions and indicators reflect the complex range of sexual behaviours that create concurrency.<sup>10</sup> However, to compare the role of concurrency across populations, a standard measure is necessary.

Currently, several countries are planning or implementing HIV-prevention strategies that specifically target the reduction of concurrent sexual partnerships.<sup>11</sup> To provide the consensus primary indicator of concurrency at the population level needed to evaluate and compare these programmes, the UNAIDS Reference Group on Estimates, Modelling, and Projections convened a meeting of international experts in sexual behaviour to agree on a definition, indicator, and method for measuring concurrent sexual partnerships.

The definition for concurrent sexual partnerships that we suggest is: overlapping sexual partnerships in which sexual intercourse with one partner occurs between two acts of intercourse with another partner. This definition embodies the generally understood meaning of concurrency,<sup>4</sup> and makes explicit the importance of sex with two different partners in temporally overlapping partnerships.

The main indicator of concurrency in a population that we recommend is the point prevalence of concurrency in the adult population, defined as the proportion of all adults in the population having more than one sexual partnership at a point in time. To calculate the indicator from data, we recommend that the point in time be 6 months before the interview (figure) so that, in most cases, it is clear whether the respondent did have sex again with a previous partner after forming a new partnership. This indicator emphasises the occurrence of sustained overlapping partnerships compared with individuals having a single long-term partnership with an occasional one-off sexual encounter, which is in line with the theoretical development of the concurrency hypothesis.<sup>1,7</sup> We recommend that any study about the frequency of concurrent partnerships in a population should include this indicator.

Other indicators that might also provide useful summaries of concurrency in a general population are the cumulative prevalence of concurrent partnerships (the proportion of all adults that have had concurrent partnerships at any point in the past year), and the proportion of individuals with multiple partnerships in the past year that had concurrent partnerships in the past year (figure). The latter indicator is intended to distinguish between the risk of concurrent sexual partnerships from

that of having multiple sexual partnerships, a well-established risk factor for HIV transmission in several settings.<sup>12</sup>

The method through which these indicators can be measured is a survey of sexual-partner histories from a random sample of adults. In these surveys, individuals are asked the following questions about each of up to their three most recent sexual partners in the past year. First, how long ago did you last have sexual intercourse with this person? Second, how long ago did you first have sexual intercourse with this person? Third, are you still having sex with this person? The phrasing of these questions should refer specifically to acts of sexual intercourse to avoid ambiguities with culturally specific notions of relationships.

Further details about the calculation and interpretation of the recommended indicators, the collection of data for measuring concurrency, and areas requiring further methodological and epidemiological research are in the meeting report.<sup>13</sup>


Considerable efforts, which include attempting to reduce concurrency, are being made to prevent HIV spread in high-prevalence settings—efforts that require monitoring and evaluation. Adoption of the common set of tools and indicators that we propose will accelerate research into understanding the relation between concurrency and HIV transmission, and enable the evaluation and promotion of successful programmes that target concurrent partnerships.

*UNAIDS Reference Group on Estimates, Modelling, and Projections: Working Group on Measuring Concurrent Sexual Partnerships*

[jeffrey.eaton08@imperial.ac.uk](mailto:jeffrey.eaton08@imperial.ac.uk)

All members of the Working Group contributed to this Comment. The meeting of the UNAIDS Reference Group at which consensus was reached for these recommendations was funded by the Joint UN Program on HIV/AIDS. We declare no conflicts of interest.

- 1 Hudson CP. Concurrent partnerships could cause AIDS epidemics. *Int J STD AIDS* 1993; 4: 249-53.
- 2 Halperin DT, Epstein H. Concurrent sexual partnerships help to explain Africa's high HIV prevalence: implications for prevention. *Lancet* 2004; 364: 4-6.
- 3 2009 AIDS Epidemic Update. Joint United Nations Programme on HIV/AIDS (UNAIDS) and World Health Organization (WHO). 2009. [http://data.unaids.org/pub/Report/2009/2009\\_epidemic\\_update\\_en.pdf](http://data.unaids.org/pub/Report/2009/2009_epidemic_update_en.pdf) (accessed Nov 24, 2009).
- 4 Shekhar N, Patts G. Measuring sexual partnership concurrency: PSI research & metrics toolkit. 2008. [http://www.psi.org/research/toolkits/measuring\\_concurrency\\_4-28.pdf](http://www.psi.org/research/toolkits/measuring_concurrency_4-28.pdf) (accessed Nov 13, 2009).
- 5 Carael M. Sexual behaviour. In: Cleland JG, Ferry B, eds. *Sexual behaviour and AIDS in the developing world*. Abingdon, UK: Taylor & Francis, 1995: 75-123.
- 6 Shelton JD. Why multiple sexual partners? *Lancet*. 2009; 374: 367-69.
- 7 Morris M, Kretzschmar M. Concurrent partnerships and the spread of HIV. *AIDS* 1997; 11: 641-48.
- 8 Mah TL, Halperin DT. Concurrent sexual partnerships and the HIV epidemics in Africa: evidence to move forward. *AIDS Behav* 2008; published online July 22. DOI:10.1007/s10461-008-9433-x.
- 9 Lurie MN, Rosenthal S. Concurrent partnerships as a driver of the HIV epidemic in sub-Saharan Africa? The evidence is limited. *AIDS Behav* 2009; published online June 2. DOI:10.1007/s10461-009-9583-5.
- 10 Gorbach PM, Stoner BP, Aral SO, Whittington WLH, Holmes KK. "It takes a village"—understanding concurrent sexual partnerships in Seattle, Washington. *Sex Transm Dis* 2002; 29: 453-62.
- 11 AIDSTAR-One. Addressing multiple and concurrent sexual partnerships in generalized HIV epidemics: report on a technical consultation in Washington, DC. October 29-30, 2008. 2009. [http://www.aidstar-one.com/sites/default/files/AIDSTAR-One\\_MCP\\_Report.pdf](http://www.aidstar-one.com/sites/default/files/AIDSTAR-One_MCP_Report.pdf) (accessed Nov 2, 2009).
- 12 Chen L, Jha P, Stirling B, et al. Sexual risk factors for HIV infection in early and advanced HIV epidemics in sub-Saharan Africa: systematic overview of 68 epidemiological studies. *PLoS One* 2007; 2: e1001.
- 13 UNAIDS. Consultation on concurrent sexual partnerships: recommendations from a meeting of the UNAIDS Reference Group on Estimates, Modelling and Projections held in Nairobi, Kenya, April 20–21st 2009. 2009. [http://www.epidem.org/Publications/Concurrency%20meeting%20recommendations\\_Final.pdf](http://www.epidem.org/Publications/Concurrency%20meeting%20recommendations_Final.pdf) (accessed Nov 21, 2009).



**Figure: Hypothetical sexual-partner histories**

Dates of first and last sex with up to three partners in past year in survey of six respondents. Point prevalence of concurrency 6 months before interview date is 33.3% (respondents 1 and 6 had concurrent partners 6 months before interview). Cumulative prevalence of concurrency is 50% (1, 3, and 6 had concurrent partnerships in past year). Proportion of individuals with multiple partnerships who had concurrent partnerships in past year is 60% (1, 3, and 6 had concurrent partnerships in past year, while 2 and 5 had multiple partnerships that were not concurrent). Respondent 5 reported both one partnership ending and another beginning 6 months before interview date. Consensus of meeting participants was that this should not be included in numerator for calculation of these indicators, as it is not possible to tell whether these partnerships are serial or concurrent.